arXiv:1204.4207v2 [hep-th] 25 Jun 2012

ON SOFT LIMITS OF INFLATIONARY CORRELATION FUNCTIONS

Valentin Assassi♠, Daniel Baumann♠, and Daniel Green♣

♠ *D.A.M.T.P., Cambridge University, Cambridge, CB3 0WA, UK*

♣ *School of Natural Sciences, Institute for Advanced Study, Princeton, NJ 08540, USA*

Abstract

Soft limits of inflationary correlation functions are both observationally relevant and theoretically robust. Various theorems can be proven about them that are insensitive to detailed model-building assumptions. In this paper, we re-derive several of these theorems in a universal way. Our method makes manifest why soft limits are such an interesting probe of the spectrum of additional light fields during inflation. We illustrate these abstract results with a detailed case study of the soft limits of quasi-single-field inflation.

Contents

1 Introduction **2**

2 Generalities: Soft Limits of Inflationary Correlation Functions **3**
- 2.1 Soft Internal Momenta 3
  - 2.1.1 Suyama-Yamaguchi Bound 3
  - 2.1.2 Interpretation 6
- 2.2 Soft External Momenta 7
  - 2.2.1 Symmetries of Inflation 7
  - 2.2.2 Consistency Relations 9
  - 2.2.3 Interpretation 10

3 Case Study: Soft Limits of Quasi-Single-Field Inflation **11**
- 3.1 Review of Quasi-Single-Field Inflation 11
- 3.2 Explicit 'in-in' Computations 13
  - 3.2.1 Power Spectrum 14
  - 3.2.2 Soft Limits: Methodology 14
  - 3.2.3 Bispectrum in the Squeezed Limit 16
  - 3.2.4 Trispectrum in the Collapsed Limit 18

4 Discussion: Non-Gaussianity as a Particle Detector **20**

A Consistency Relations at Finite Momentum **22**

1 Introduction

Soft limits of primordial correlation functions are of special phenomenological relevance, since many observational probes of non-Gaussianity—such as scale-dependent halo bias [1] and CMB μ-distortions [2]—involve a separation of scales. At the same time, soft limits are theoretically clean. Focusing on soft limits allows one to prove a number of interesting theorems that are insensitive to detailed model-building assumptions, but are sensitive to the number of light degrees of freedom during inflation. Moreover, as we will explain in this paper, the momentum scaling in the soft limits is sensitive to details of the spectrum of particles during inflation.

There are two classes of soft limits:

1. The 'squeezed limit' of an N-point function refers to taking one external momentum to be smaller than all others (see Fig. 1). Maldacena's consistency relation for single-field inflation [3] (see also [4–6]) relates the squeezed limit of the three-point function of curvature perturbations ζ to the variation of the two-point function under dilation. Similar results have subsequently been derived for N-point functions [7–10]. These results are explained by realizing that the curvature perturbation ζ non-linearly realizes a dilation symmetry in any FRW background [11] (see also [12–14]). Any violation of the consistency relations would rule out *all* single-field models and point to additional degrees of freedom during inflation.



Figure 1: Soft external momentum or the 'squeezed limit'.

2. The 'collapsed limit' of an N-point function takes one internal momentum (i.e. the sum of M of the N external momenta) to be smaller than all external momenta (see Fig. 2). The Suyama-Yamaguchi inequality [15] (see also [16, 17, 20]) puts a bound on the collapsed limit of the four-point function in terms of the squeezed limit of the three-point function. At tree level, the inequality is saturated if a 'single source' creates the primordial curvature



Figure 2: Soft internal momentum or the 'collapsed limit'.

> perturbation and its non-Gaussianity [21]. Collapsed limits are therefore probes of 'multiple source' (or 'hidden sector') non-Gaussianities. An interesting observational signature of this effect is stochasticity in the scale-dependent halo bias [22] (see [23] for another possible signature).

A boost in the collapsed limits arises naturally in models of quasi-single field inflation (QSFI) [24]. In these models a light inflaton field mixes with an additional scalar field (the 'isocurvaton') with mass of order the Hubble scale H.[1] This mixing allows large non-Gaussianities in the isocurvaton sector to be communicated to the observable sector. The time dependence of the massive field on superhorizon scales leads to a characteristic scaling behavior of N-point functions in their soft limits [25]. Moreover, the perturbative coupling between the observable sector and the isocurvature sector leads to the boost in the collapsed limit relative to the squeezed limit.

The outline of the paper is as follows: In Section 2, we present new proofs of various important theorems [3, 15] concerning the soft limits of primordial correlation functions. Although these results aren't new, some of our methods and perspectives may be. In particular, we make a bit more precise the analogy between soft limits of inflationary correlation functions and soft pion limits of QCD [26–31]. In Appendix A, we present a number of technical details and generalizations related to the theorems of Section 2. In Section 3, we compute the soft limits of QSFI explicitly in terms of the fundamental parameters of the theory. We conclude in Section 4 with a discussion of the prospects of using non-Gaussianity as a particle detector.

2 Generalities: Soft Limits of Inflationary Correlation Functions

We begin with proofs of various important theorems [3, 15] concerning the soft limits of inflationary correlation functions.

2.1 Soft Internal Momenta

We first consider the collapsed limit of the four-point function, i.e. we let the internal momentum of the four-point function become soft, while keeping all external momenta hard. We will show that the four-point function in this limit is bounded from below by the squeezed limit of the corresponding three-point function. Our proof will make the role of additional degrees of freedom manifest. It will also make obvious how the proof generalizes to higher N-point functions.

2.1.1 Suyama-Yamaguchi Bound

Definitions.—We characterize the soft limits of the three- and four-point functions by the following two quantities

$$\hat{f}_{\rm NL} \equiv \frac{5}{12} \lim_{k_1\to 0} \frac{\langle \zeta_{\vec{k}_1}\zeta_{\vec{k}_2}\zeta_{\vec{k}_3}\rangle'}{P_1 P_2} \ , \tag{2.1}$$

$$\hat{\tau}_{\rm NL} \equiv \frac{1}{4} \lim_{k_{12}\to 0} \frac{\langle \zeta_{\vec{k}_1}\zeta_{\vec{k}_2}\zeta_{\vec{k}_3}\zeta_{\vec{k}_4}\rangle'}{P_1 P_3 P_{12}} \ , \tag{2.2}$$

---

[1] Such a spectrum of particle masses arises generically in supersymmetric theories of inflation [25].

where $\langle \cdot \rangle'$ denotes the expectation value without the factor $(2\pi)^3 \delta(\sum \vec{k}_i)$. Moreover, we have defined $P_i \equiv P_\zeta(k_i) = \langle \zeta_{\vec{k}_i} \zeta_{\vec{k}_j} \rangle'$ and $k_{12} \equiv |\vec{k}_1 + \vec{k}_2|$. All correlation functions are to be understood as 'connected' correlation functions. For local non-Gaussianity, the definitions (2.1) and (2.2) reduce to the familiar (momentum-independent) definitions of $f_{\rm NL}$ [32] and $\tau_{\rm NL}$ [21]. More generally, however, $\hat{f}_{\rm NL}$ and $\hat{\tau}_{\rm NL}$ can be momentum dependent—e.g. in §3, we will find $\hat{f}_{\rm NL}(k_1) \propto k_1^{3-\alpha}$ and $\hat{\tau}_{\rm NL}(k_{12}) \propto k_{12}^{6-2\alpha}$, with $3 \geq \alpha \geq \frac{3}{2}$. We will now give a proof of the bound $\hat{\tau}_{\rm NL} \geq (\frac{6}{5}\hat{f}_{\rm NL})^2$ [15, 17]. In fact, we will give a proof of a slightly more general inequality, which includes $\hat{\tau}_{\rm NL} \geq (\frac{6}{5}\hat{f}_{\rm NL})^2$ as a special case.

Collapsed four-point function.—We start by considering the following four-point function,[2]

$$\langle \zeta^2(\vec{x}, t_\star) \zeta^2(\vec{0}, t_\star) \rangle \equiv \langle \Omega | \zeta^2(\vec{x}, t_\star) \zeta^2(\vec{0}, t_\star) | \Omega \rangle \ . \tag{2.3}$$

where $t_\star$ is some time at which ζ has become frozen and $|\Omega\rangle$ is the vacuum[3] of the interacting theory. Next, we insert a complete set of momentum eigenstates $|n_{\vec{q}}\rangle$ in the correlation function,

$$\langle \zeta^2(\vec{x}, t_\star) \zeta^2(\vec{0}, t_\star) \rangle = \sum_{n, \vec{q}} \langle \zeta^2(\vec{x}, t_\star) | n_{\vec{q}} \rangle \langle n_{\vec{q}} | \zeta^2(\vec{0}, t_\star) \rangle \ , \tag{2.4}$$

where $\sum_{n,\vec{q}}(\cdot) \equiv \sum_n \int_{\vec{q}} (\cdot) = \sum_n \int \frac{{\rm d}^3 \vec{q}}{(2\pi)^3} (\cdot)$. We are working in the Heisenberg picture and therefore the states are formally[4] independent of time. Here, $\vec{q}$ is the total momentum of a state and n is a label for individual states (e.g. 'particle number'[5] states or different particle species) within the momentum eigenspace. In the following, we will drop the time label $t_\star$. Using $\langle \zeta^2(\vec{x}) | n_{\vec{q}} \rangle = e^{i \vec{q} \cdot \vec{x}} \langle \zeta^2(\vec{0}) | n_{\vec{q}} \rangle$, we can write eq. (2.4) as

$$\langle \zeta^2(\vec{x}) \zeta^2(\vec{0}) \rangle_{\vec{k}} \equiv \int {\rm d}^3 x \, e^{-i \vec{k} \cdot \vec{x}} \langle \zeta^2(\vec{x}) \zeta^2(\vec{0}) \rangle = \sum_n |\langle n_{\vec{k}} | \zeta^2(\vec{0}) \rangle|^2 \ . \tag{2.5}$$

Notice that the r.h.s. is written as a sum over positive definite terms. If we can show that one of the contributions is the square of the squeezed limit of the three-point function of ζ, then we can conclude that the collapsed four-point function—the l.h.s. of (2.5)—is always greater or equal.

Squeezed three-point function.—To complete the proof, we insert a complete set of states in the three-point function $\langle \zeta(\vec{x}) \zeta^2(\vec{0}) \rangle$, i.e.

$$\langle \zeta_{\vec{k}} | \zeta^2(\vec{0}) \rangle = \int {\rm d}^3 x \, e^{+i \vec{k} \cdot \vec{x}} \langle \zeta(\vec{x}) \zeta^2(\vec{0}) \rangle = \sum_{n, \vec{q}} \langle \zeta_{\vec{k}} | n_{\vec{q}} \rangle \langle n_{\vec{q}} | \zeta^2(\vec{0}) \rangle \ , \tag{2.6}$$

where $\langle \zeta_{\vec{k}} | = | \zeta_{\vec{k}} \rangle^\dagger = \langle \Omega | \zeta_{-\vec{k}}$. It is always possible to choose the basis states $|n_{\vec{q}}\rangle$, such that

$$\langle \zeta_{\vec{k}} | n_{\vec{q}} \rangle = (2\pi)^3 \delta(\vec{k} - \vec{q}) \, \delta_{1n} \, P_\zeta^{1/2}(k) \ . \tag{2.7}$$

Up to an arbitrary phase, this implies $|1_{\vec{k}}\rangle = P_\zeta^{-1/2}(k) \, |\zeta_{\vec{k}}\rangle$. In an abuse of the standard particle physics terminology, we will call $|1_{\vec{k}}\rangle$ the 'one-zeta' state. All other states, which we will denote

---

[2] Our conclusions will be unchanged if we replace $\zeta^2(\vec{x}, t_\star) \to \zeta(\vec{x}, t_\star)\zeta(\vec{x} + \vec{\varepsilon}, t_\star)$ and $\zeta^2(\vec{0}, t_\star) \to \zeta(\vec{0}, t_\star)\zeta(\vec{\varepsilon}, t_\star)$, which may be useful in order to avoid divergences (or contact terms) when the operators are at coincident points.

[3] It is worth remarking that our results hold even if $|\Omega\rangle$ is not the vacuum state, as long as the state is invariant

$\|\Omega\rangle$	vacuum
$\|1_{\vec{k}}\rangle \propto \|\zeta_{\vec{k}}\rangle$	"one-zeta"
$\|\tilde{n}_{\vec{k}}\rangle \perp \|\zeta_{\vec{k}}\rangle$	"everything else"

Table 1: Spectrum of states and their meanings.

by $|\tilde{n}_{\vec{k}}\rangle$, are orthogonal to $|\zeta_{\vec{k}}\rangle$. They may be associated with 'single-particle' states corresponding to additional degrees of freedom or simply 'multi-zeta' states, i.e. states created with higher powers of the operator ζ. Substituting (2.7) into (2.6), we get

$$\langle \zeta_{\vec{k}} | \zeta^2(\vec{0})\rangle = \langle 1_{\vec{k}} | \zeta^2(\vec{0})\rangle \, P_\zeta^{1/2}(k) \ . \tag{2.8}$$

Bound on the collapsed four-point function.—We can therefore write eq. (2.5) as

$$\boxed{\langle \zeta^2(\vec{x})\zeta^2(\vec{0})\rangle_{\vec{k}} \ = \ \frac{|\langle \zeta_{\vec{k}} | \zeta^2(\vec{0})\rangle|^2}{P_\zeta(k)} \ + \ \sum_{\tilde{n}} |\langle \tilde{n}_{\vec{k}} | \zeta^2(\vec{0})\rangle|^2} \ . \tag{2.9}$$

Since $\sum_{\tilde{n}} |\langle \tilde{n}_{\vec{k}} | \zeta^2(\vec{0})\rangle|^2 \ \geq \ 0$, we conclude that

$$\langle \zeta^2(\vec{x})\zeta^2(\vec{0})\rangle_{\vec{k}} \ \geq \ \frac{|\langle \zeta_{\vec{k}} | \zeta^2(\vec{0})\rangle|^2}{P_\zeta(k)} \ . \tag{2.10}$$

If we take the soft limit $k \to 0$, we get a relation between the collapsed four-point function and the squeezed three-point function. Moreover, from our derivation it is easy to see how the result can be generalized to give a relation between the soft limits of $2N$-point functions and the squeezed limits of $(N+1)$-point functions,

$$\langle \zeta^N(\vec{x})\zeta^N(\vec{0})\rangle_{\vec{k}} \ = \ \frac{|\langle \zeta_{\vec{k}} | \zeta^N(\vec{0})\rangle|^2}{P_\zeta(k)} \ + \ \sum_{\tilde{n}} |\langle \tilde{n}_{\vec{k}} | \zeta^N(\vec{0})\rangle|^2 \ . \tag{2.11}$$

Suyama-Yamaguchi bound.—Eq. (2.10) can be written as

$$\int_{\vec{q}_1}\int_{\vec{q}_2} \langle \zeta_{\vec{q}_1}\zeta_{\vec{k}-\vec{q}_1}\zeta_{\vec{q}_2}\zeta_{-\vec{q}_2-\vec{k}}\rangle' \ \geq \ \frac{\left|\int_{\vec{q}_1}\langle \zeta_{\vec{k}}\zeta_{\vec{q}_1}\zeta_{-\vec{q}_1-\vec{k}}\rangle'\right|^2}{P_\zeta(k)} \ , \tag{2.12}$$

which, in the limit $k \to 0$, becomes

$$\int_{\vec{q}_1}\int_{\vec{q}_2} \hat{\tau}_{\rm NL}\, P_\zeta(q_1)P_\zeta(q_2) \ \geq \ \int_{\vec{q}_1}\int_{\vec{q}_2} (\tfrac{6}{5}\hat{f}_{\rm NL})^2\, P_\zeta(q_1)P_\zeta(q_2) \ . \tag{2.13}$$

---

under spatial translations and rotations.

[4] In practice, we define the Hilbert space perturbatively using the free theory. The states then become implicitly time dependent through the projection onto the interacting vacuum.

[5] Of course, the notion of 'particle states' is subtle in time-dependent cosmological backgrounds without asymptotically free states. None of those subtleties will affect our arguments, so we will, for convenience, sometimes (ab)use the particle physics terminology.

So far this is completely general. To get the specific form of the Suyama-Yamaguchi bound [15] one has to make a further assumption. It is often the case that $\hat{\tau}_{\rm NL}$ and $(\hat{f}_{\rm NL})^2$ are either momentum independent or have the same momentum dependence (so that we can still extract a momentum-independent amplitude). We then get

$$\boxed{\hat{\tau}_{\rm NL} \geq (\tfrac{6}{5}\hat{f}_{\rm NL})^2}\ . \tag{2.14}$$

2.1.2 Interpretation

What is the physical meaning of the extra contributions in eq. (2.9)? It is well-known [15, 21] that curvature fluctuations arising from a 'single source' (which may or may not be the inflaton) saturate the Suyama-Yamaguchi inequality (at least at tree level). In this case, the squeezed three-point function of ζ completely determines the collapsed four-point function—i.e. the extra contributions in (2.9) are subdominant, in the sense that

$$\lim_{k\to 0} \frac{\langle \tilde{n}_{\vec{k}} | \zeta^2 \rangle}{\langle 1_{\vec{k}} | \zeta^2 \rangle} = 0 \qquad \text{(single source at tree level)}\ . \tag{2.15}$$

Non-negligible $\langle \tilde{n}_{\vec{k}\to 0} | \zeta^2 \rangle$ can have the following origins:

1. *Extra fields.* A straightforward way to have contributions from extra states is to have multiple sources for the primordial fluctuations. This can lead to a significant boost of the collapsed four-point function relative to the square of the squeezed three-point function, $\hat{\tau}_{\rm NL} \gg (\frac{6}{5}\hat{f}_{\rm NL})^2$. As a measure of such 'multi-source' non-Gaussianity we can define

$$X \equiv \lim_{k\to 0} \left[\frac{\langle \zeta^2(\vec{x})\zeta^2(\vec{0})\rangle_{\vec{k}}\, P_\zeta(k)}{|\langle \zeta_{\vec{k}} | \zeta^2(\vec{0})\rangle|^2} - 1 \right] = \lim_{k\to 0} \sum_{\tilde{n}} \frac{|\langle \tilde{n}_{\vec{k}} | \zeta^2(\vec{0})\rangle|^2}{|\langle 1_{\vec{k}} | \zeta^2(\vec{0})\rangle|^2}\ . \tag{2.16}$$

   We will see an explicit example of this situation in Section 3.

2. *Gravitons.* Strictly speaking, even single-source inflation always has additional states corresponding to the graviton, i.e. tensor metric perturbations. In fact, in single-field *slow-roll* inflation these states give the dominant contribution to the collapsed four-point function, $\hat{\tau}_{\rm NL} \sim \mathcal{O}(\epsilon)$ [10], where $\epsilon \equiv -\dot{H}/H^2 < 1$. This is parametrically larger than $\hat{f}^2_{\rm NL} \sim \mathcal{O}(\epsilon^2)$ [3], although both contributions are too small to be observable. Being interested in observable signatures, we will typically suppress the graviton contributions in our results. It would be easy to include their effects as $\mathcal{O}(\epsilon)$ corrections.

3. *Excited initial states.* The effects of excited initial states on primordial non-Gaussianity have been explored in [35–40]. In particular, it was pointed out [37] that non-trivial contributions to soft limits are possible. It is conceivable that these effects are captured by $\langle \tilde{n}_{\vec{k}\to 0} | \zeta^2 \rangle$ in our formalism, but we leave a detailed analysis for the future [41].

4. *Loops.* Even in single-source inflation we can (formally) have $\hat{\tau}_{\rm NL} > (\frac{6}{5}\hat{f}_{\rm NL})^2$ from loops [18–20]. In single-clock inflation [33], these loop contributions aren't physical and can be removed by the right choice of physical coordinates [34]. In more general single-source scenarios, loops may contribute non-trivially. Having said that, in this paper, we won't consider these loop effects, but instead focus on tree-level contributions.

2.2 Soft External Momenta

We now use similar ideas to derive the famous consistency relations for correlation functions with soft external momenta [3, 4]. Our approach will draw inspiration from methods used in proving the low-energy theorems of theories with spontaneous symmetry breaking [29, 30], such as the classic soft pion theorems of QCD [26–28]. This connection was emphasized recently in [11]. The proofs in this section will be restricted to the limit $k \to 0$, but we will show in Appendix A how they can be generalized to finite k.

2.2.1 Symmetries of Inflation

The consistency relations arise from the symmetries of inflation, which we therefore quickly review (for further discussion see [11, 42, 43]).



Figure 3: Conformal diagram of de Sitter space. The inflationary background picks out a preferred time-slicing.

Inflation is characterized by a quasi-de Sitter background

$$\mathrm{d}s^2 \simeq -\mathrm{d}t^2 + e^{2Ht}\mathrm{d}\vec{x}^2 \ . \tag{2.17}$$

The isometry group of de Sitter space, $SO(4,1)$, contains 6 translations and rotations, 1 dilation

$$Ht \ \mapsto \ Ht - \lambda \ , \qquad \vec{x} \ \mapsto \ (1+\lambda)\vec{x} \ , \tag{2.18}$$

and 3 special conformal transformations (SCTs)

$$Ht \ \mapsto \ Ht - 2\vec{b}\cdot\vec{x} \ , \qquad \vec{x} \ \mapsto \ \vec{x} + \vec{b}(H^{-2}e^{-2Ht} - \vec{x}^2) + 2(\vec{b}\cdot\vec{x})\vec{x} \ , \tag{2.19}$$

where λ and $\vec{b}$ are infinitesimal parameters. At late times, $t \to \infty$, these isometries act as conformal transformations on the spatial boundary $\mathcal{I}^+$:

$$\vec{x} \ \mapsto \ (1+\lambda)\vec{x} \ , \tag{2.20}$$

$$\vec{x} \ \mapsto \ \vec{x} - \vec{b}\vec{x}^2 + 2(\vec{b}\cdot\vec{x})\vec{x} \ . \tag{2.21}$$

The time-dependence of the inflationary background $\phi(t)$ spontaneously breaks the symmetry under dilations and special conformal transformations. Inflationary correlations are therefore, in general, not invariant under the conformal transformations.[6] However, since the symmetry is non-linearly realized, we still expect non-trivial relations between different N-point functions.

As explained nicely in recent work by Creminelli, Noreña, and Simonović [12] and Hinterbichler, Hui and Khoury [11], the symmetries in eqs. (2.20) and (2.21), in fact, have significance beyond quasi-de Sitter backgrounds. They are symmetries of adiabatic modes in any FRW background.[7] The Goldstone boson associated with the spontaneous breaking of the symmetry is the curvature perturbation[8] ζ, which in co-moving gauge is defined by [47, 48]

$$g_{ij}(\vec{x},t) = a^2(t)e^{2\zeta(\vec{x},t)}\delta_{ij} \ . \tag{2.23}$$

As we will see, the squeezed limit of N-point functions of ζ are related to the variation of $(N-1)$-point functions under dilations and special conformation transformations. This is a consequence of the conformal transformations being non-linearly realized.

We now show how these considerations can be incorporated into our formalism. For concreteness, we focus on the dilation symmetry[9] in co-moving coordinates

$$\vec{x} \ \mapsto \ (1+\lambda)\vec{x} \ , \tag{2.24}$$

$$\zeta \ \mapsto \ \zeta + \lambda(1+\vec{x}\cdot\partial_{\vec{x}}\zeta) \ . \tag{2.25}$$

This symmetry is a large[10] diffeomorphism that is non-linearly realized by the "gauge-invariant" perturbation ζ. It should not be confused with an isometry of FRW (which is linearly realized) or a residual[11] gauge transformation. During inflation, only time diffeomorphisms are broken by the background [33]. Therefore, this large spatial diffeomorphism will act linearly on all the non-gravitational degrees of freedom. Associated with the dilation symmetry of the action is a conserved current J_d^μ and a charge $Q_d \equiv \int \mathrm{d}^3x\, J_d^0(\vec{x},t)$. Although the charge does not exist after symmetry breaking (due to an IR divergence), local commutators are still well defined, e.g.

$$\delta_d\zeta \equiv i[Q_d,\zeta] = -1 - \vec{x}\cdot\partial_{\vec{x}}\zeta \ . \tag{2.26}$$

For future reference, we will need the matrix element $\langle\Omega|Q_d\zeta_{\vec{q}}|\Omega\rangle \equiv \langle Q_d|\zeta_{\vec{q}}\rangle$. Invariance of the vacuum under spatial translations and rotations, fixes the form of this matrix element to be

$$\langle Q_d|\zeta_{\vec{q}}\rangle = i f(q)\cdot(2\pi)^3\delta(\vec{q}) \ , \tag{2.27}$$

---

[6] The symmetries are still explicit for fields that don't couple to the inflaton, such as tensor metric fluctuations [42] or scalar spectator fields [43, 44]. Even for inflaton fluctuations, the symmetries are restored in the so-called decoupling limit, where $M_{\rm pl}\to\infty$, $\dot{H}\to 0$, with $M_{\rm pl}^2\dot{H}\to$ *const*.

[7] Adiabatic modes are defined locally as coordinate redefinitions of the unperturbed FRW solution. Coordinate transformations of the form (2.20) and (2.21) with time-dependent parameters $\lambda(t)$ and $\vec{b}(t)$ induce the following co-moving curvature perturbation

$$\zeta = \lambda(t) + 2\vec{b}(t)\cdot\vec{x} \ . \tag{2.22}$$

For *constant* transformation parameters λ and $\vec{b}$ this gauge mode is the $k\to 0$ limit of a physical solution [11, 12], a la Weinberg [45], with suitable fall-off at spatial infinity.

[8] For spontaneously broken 'spacetime' symmetries the number of Goldstone bosons does not have to equal the number of broken generators [46].

[9] It is our understanding that a complete treatment will appear in [13, 14] (see also [12]).

[10] A "large diffeomorphism" is one that does not vanish at infinity. These transformations are distinct from the gauge redundancies of the theory, namely local diffeomorphisms that vanish at infinity.

[11] There is a residual gauge transformation in "ζ-gauge", eq. (2.23), that does not act on ζ [49].

where f is a function that depends only on the magnitude of the 3-momentum $q \equiv |\vec{q}|$. In order for this to be consistent with the commutator (2.26), we require $\mathrm{Re}[f(q)] = \frac{1}{2}$. The imaginary part of $f(q)$ won't contribute in what we are about to say, so we can set it to zero. We then have

$$\langle Q_d|\zeta_{\vec{q}}\rangle = \tfrac{i}{2} \cdot (2\pi)^3 \delta(\vec{q}) \ . \tag{2.28}$$

2.2.2 Consistency Relations

We now use these results to give a quick proof of the 'Adler zero' [26] for the squeezed limit of the three-point function in single-field inflation [3, 4]. We will also see how extra fields have to enter in order to produce a non-trivial signal in the squeezed limit.

Associated with the conserved current J^μ_d is the following Ward identity [50]

$$\begin{aligned} i\,\partial^{(x)}_\mu \langle J^\mu_d(\vec{x},t)\zeta(\vec{y},t_\star)\zeta(\vec{z},t_\star,)\rangle \ &= \ \delta(t-t_\star)\delta(\vec{x}-\vec{y})\,\langle \delta_d\zeta(\vec{y},t_\star)\,\zeta(\vec{z},t_\star)\rangle \\ &\quad + \delta(t-t_\star)\delta(\vec{x}-\vec{z})\,\langle \zeta(\vec{y},t_\star)\,\delta_d\zeta(\vec{z},t_\star)\rangle \ , \end{aligned} \tag{2.29}$$

where for clarity we have this time shown all the time dependences explicitly. If we integrate[12] eq. (2.29) over $\int_{t_\star - i\epsilon}^{t_\star + i\epsilon} \mathrm{d}t \int \mathrm{d}^3x$, we arrive at

$$\langle [Q_d, \zeta_{\vec{k}_1}\zeta_{\vec{k}_2}]\rangle \ = \ \langle [Q_d, \zeta_{\vec{k}_1}]\zeta_{\vec{k}_2}\rangle + \langle \zeta_{\vec{k}_1}[Q_d, \zeta_{\vec{k}_2}]\rangle \ , \tag{2.30}$$

where we have reverted to dropping the common time coordinate $t_\star$. In going from (2.29) to (2.30), we have dropped the integral over $\partial_i J^i_d$ because it only receives contributions from spatial infinity (see Appendix A). With the help of (2.26), we can express the r.h.s. of (2.30) as

$$\langle [Q_d, \zeta_{\vec{k}_1}\zeta_{\vec{k}_2}]\rangle = -i(3 + \vec{k}_1 \cdot \partial_{\vec{k}_1} + \vec{k}_2 \cdot \partial_{\vec{k}_2})\langle \zeta_{\vec{k}_1}\zeta_{\vec{k}_2}\rangle'\,(2\pi)^3\delta(\vec{k}_1 + \vec{k}_2) \ . \tag{2.31}$$

Here and in the following, we only keep terms that contribute to the connected part of the correlation functions. Next, we insert a complete set of states in the l.h.s. of (2.30),

$$\langle [Q_d, \zeta_{\vec{k}_1}\zeta_{\vec{k}_2}]\rangle \ = \ \sum_{n,\vec{q}} \Big[\langle Q_d|n_{\vec{q}}\rangle\langle n_{\vec{q}}|\zeta_{\vec{k}_1}\zeta_{\vec{k}_2}\rangle - \langle \zeta_{\vec{k}_1}\zeta_{\vec{k}_2}|n_{\vec{q}}\rangle\langle n_{\vec{q}}|Q_d\rangle\Big] \ , \tag{2.32}$$

where

$$\langle Q_d|n_{\vec{q}}\rangle \equiv \tfrac{1}{2}c_n(q) \cdot (2\pi)^3\delta(\vec{q}) \ . \tag{2.33}$$

Since $c_0 = \langle Q_d\rangle' \equiv \langle \Omega|Q_d|\Omega\rangle' \in \mathbb{R}$, the vacuum insertion doesn't contribute. We separate the 'one-zeta' state, with $c_1(q) = \frac{i}{2}P_\zeta^{-1/2}(q)$ (up to a removable phase), from the rest[13]

$$\langle [Q_d, \zeta_{\vec{k}_1}\zeta_{\vec{k}_2}]\rangle \ = \ \lim_{q\to 0}\left[\frac{i}{P_\zeta(q)}\langle \zeta_{\vec{q}}|\zeta_{\vec{k}_1}\zeta_{\vec{k}_2}\rangle' + i\sum_{\tilde{n}} \mathrm{Im}\Big[c_{\tilde{n}}(q)\langle \tilde{n}_{\vec{q}}|\zeta_{\vec{k}_1}\zeta_{\vec{k}_2}\rangle'\Big]\right](2\pi)^3\delta(\vec{k}_1 + \vec{k}_2) \ . \tag{2.34}$$

---

[12] The integral over time is extended in the imaginary direction because our correlation functions are "time ordered" with respect to the integration contour in the complex plane (see Appendix A for further details).

[13] The functions $|c_{\tilde{n}}(q)|^2$ in eq. (2.34) are analogous to the spectral density functions $\rho(q^2)$ in the Källén-Lehmann representation of correlation functions [29]. In principle, one could determine the properties of $|c_{\tilde{n}}(q)|^2$ from $\langle J^\mu_d(\vec{x})J^\nu_d(\vec{0})\rangle$.

In deriving eq. (2.34) we have used the fact that

$$\langle\zeta_{\vec{k}_1}\zeta_{\vec{k}_2}|\tilde{n}_{\vec{q}}\rangle = \langle\tilde{n}_{\vec{q}}|\zeta_{-\vec{k}_1}\zeta_{-\vec{k}_2}\rangle^* = \langle\tilde{n}_{-\vec{q}}|\zeta_{\vec{k}_1}\zeta_{\vec{k}_2}\rangle^* \ , \tag{2.35}$$

where the second equality comes from the invariance of correlation functions under parity. Comparing eqs. (2.31) and (2.34), we get

$$\boxed{\frac{\langle\zeta_{\vec{q}}|\zeta_{\vec{k}_1}\zeta_{\vec{k}_2}\rangle'}{P_\zeta(q)} \xrightarrow{q\to 0} -\Big[3+\vec{k}_1\cdot\partial_{\vec{k}_1}+\vec{k}_2\cdot\partial_{\vec{k}_2}\Big]\langle\zeta_{\vec{k}_1}\zeta_{\vec{k}_2}\rangle' - \sum_{\tilde{n}}\mathrm{Im}\Big[c_{\tilde{n}}(q)\langle\tilde{n}_{\vec{q}}|\zeta_{\vec{k}_1}\zeta_{\vec{k}_2}\rangle'\Big]} \ . \tag{2.36}$$

This result generalizes straightforwardly to the squeezed limits of higher N-point functions (see Fig. 1)

$$\begin{aligned}\frac{\langle\zeta_{\vec{q}}|\zeta_{\vec{k}_1}\cdots\zeta_{\vec{k}_N}\rangle'}{P_\zeta(q)} \xrightarrow{q\to 0} & -\Big[3(N-1)+\sum_{i=1}^N\vec{k}_i\cdot\partial_{\vec{k}_i}\Big]\langle\zeta_{\vec{k}_1}\cdots\zeta_{\vec{k}_N}\rangle' \\ & -\sum_{\tilde{n}}\mathrm{Im}\Big[c_{\tilde{n}}(q)\langle\tilde{n}_{\vec{q}}|\zeta_{\vec{k}_1}\cdots\zeta_{\vec{k}_N}\rangle'\Big] \ .\end{aligned} \tag{2.37}$$

Moreover, it is easy to see how this result implies consistency relations for soft internal momenta [51] (see Fig. 2),

$$\begin{aligned}\frac{\langle(\zeta_{\vec{k}_1}\ldots\zeta_{\vec{k}_M})(\zeta_{\vec{k}_{M+1}}\ldots\zeta_{\vec{k}_N})\rangle'}{P_\zeta(q)} \xrightarrow{q\to 0} & \frac{\langle\zeta_{\vec{k}_1}\ldots\zeta_{\vec{k}_N}|\zeta_{\vec{q}}\rangle'}{P_\zeta(q)}\frac{\langle\zeta_{\vec{q}}|\zeta_{\vec{k}_{M+1}}\ldots\zeta_{\vec{k}_N}\rangle'}{P_\zeta(q)} \\ & -\frac{1}{P_\zeta(q)}\sum_{\tilde{n}}\langle\zeta_{\vec{k}_1}\ldots\zeta_{\vec{k}_M}|\tilde{n}_{\vec{q}}\rangle'\langle\tilde{n}_{\vec{q}}|\zeta_{\vec{k}_{M+1}}\ldots\zeta_{\vec{k}_N}\rangle' \ ,\end{aligned} \tag{2.38}$$

where $\vec{q}\equiv\sum_{i=1}^M\vec{k}_i$. Again, this was derived by inserting a complete set of states and separating the 'one-*zeta*' state $|1_{\vec{q}}\rangle = P_\zeta^{-1/2}(q)|\zeta_{\vec{q}}\rangle$ from all states orthogonal to it, $|\tilde{n}_{\vec{q}}\rangle$.

2.2.3 Interpretation

There are two important consequences of the $q\to 0$ limit. First, in this limit causality explains the absence of a potential boundary term arising from the $\partial_i J^i_d$ term in (2.29) (see Appendix A for the complete argument). Second, in this limit the 'multi-particle' states among $|\tilde{n}_{\vec{q}}\rangle$ are not expected to contribute the leading divergent term, so that we can drop them on the r.h.s. of (2.36). It is straightforward to check this in most examples, however, a general proof[14] is beyond the scope of this paper.

In the special case of single-field inflation, eq. (2.36) therefore reduces to the famous single-field consistency relation [3][15]

$$\frac{\langle\zeta_{\vec{q}}|\zeta_{\vec{k}_1}\zeta_{\vec{k}_2}\rangle'}{P_\zeta(q)} \xrightarrow{q\to 0} -\Big[3+\vec{k}_1\cdot\partial_{\vec{k}_1}+\vec{k}_2\cdot\partial_{\vec{k}_2}\Big]\langle\zeta_{\vec{k}_1}\zeta_{\vec{k}_2}\rangle' \ = \ (1-n_s)P_\zeta(k_1) \ . \tag{2.39}$$

---

[14] Even in the context of S-matrix elements for the scattering of Goldstone bosons, there are known exceptions to this statement [29] (§19.2).

[15] Excited initial states may contribute corrections at finite q, or when the number of e-folds is taken to infinity before taking $q\to 0$ (see e.g. [36–40]). This is related to the observation that $q\ll k_{1,2}$ is not a sufficient condition for satisfying the consistency relation (see e.g. [6, 54, 55]). Understanding these finite q effects can be important given that observations will only probe a finite range of momenta.

For scale-invariant fluctuations,[16] i.e. $\langle \zeta_{\vec{k}_1} \zeta_{\vec{k}_2} \rangle' = (k_1 k_2)^{-3/2}$ or $n_s = 1$, the r.h.s. of (2.39) is zero and the squeezed limit of the three-point function vanishes. This is analogous to the Adler zero of soft pion physics [26] (see also [29, 30]).

In multi-field inflation, we can get contributions to the squeezed limit, even for scale-invariant fluctuations,

$$\frac{\langle \zeta_{\vec{q}} | \zeta_{\vec{k}_1} \zeta_{\vec{k}_2} \rangle'}{P_\zeta(q)} \xrightarrow{q \to 0} -\sum_{\tilde{n}} \mathrm{Im}\Big[c_{\tilde{n}}(q) \langle \tilde{n}_{\vec{q}} | \zeta_{\vec{k}_1} \zeta_{\vec{k}_2} \rangle' \Big] . \tag{2.40}$$

This expression doesn't just tell us that we need extra fields to generate a squeezed limit, but also how the extra fields must enter. We will study an explicit example of this situation in Section 3.

3 Case Study: Soft Limits of Quasi-Single-Field Inflation

In this section, we will illustrate some of the abstract results of the previous section with the concrete example of quasi-single-field inflation (QSFI) [24]. This is a particularly interesting case study, since QSFI exhibits non-trivial behaviour in its soft limits:

i) the soft limits of N-point functions have characteristic momentum scalings that depend on the mass of the isocurvaton field;

ii) the collapsed limits of $2N$-point functions are naturally boosted.

In §3.1, we explain these features of QSFI qualitatively. In §3.2, we compute the soft limits of the three- and four-point functions explicitly.

3.1 Review of Quasi-Single-Field Inflation



Figure 4: Particle spectrum of quasi-single-field inflation.

Dynamics.—QSFI couples a massive 'isocurvaton' field σ to an 'adiabatic' perturbation θ (see Fig. 4). The dynamics of these fields is governed at leading order by the following free field Hamiltonian (density)

$$\mathcal{H}_0 = \tfrac{1}{2}(\partial_\mu \theta)^2 + \tfrac{1}{2}(\partial_\mu \sigma)^2 + \tfrac{1}{2} m^2 \sigma^2 , \tag{3.1}$$

---

[16] Scale-invariance is natural if the theory has an internal shift symmetry, $\phi \to \phi + c$. Both this symmetry and the 'spacetime' dilation symmetry of de Sitter space are spontaneously broken by the time-dependent inflationary background, $\phi(t)$. However, a linear combination of shift and spacetime dilation is preserved [52]. At late times, this diagonal symmetry becomes the spatial dilation symmetry that we discuss in this section. This explains the scale-invariance of the two-point function [53]. Softly breaking the shift symmetry allows for weakly scale-dependent fluctuation spectra.

where $m \sim H$. At linear order, the adiabatic perturbation θ and the curvature perturbation ζ are related by

$$\theta = -\sqrt{2\epsilon}\, M_{\rm pl}\, \zeta \ , \tag{3.2}$$

where $\epsilon \equiv -\dot{H}/H^2$. A quadratic mixing term, e.g. $\mathcal{H}_{\rm mix} = -\rho\dot{\theta}\sigma$, converts σ into θ (or ζ). This can communicate large interactions in the isocurvaton sector to the observable sector. Our basic example will be the cubic interaction $\mathcal{H}_3 = \mu\sigma^3$. Following [24], we treat $\mathcal{H}_{\rm mix}$ as part of the interaction Hamiltonian

$$\mathcal{H}_{\rm int} = -\rho\dot{\theta}\sigma + \mu\sigma^3 \ . \tag{3.3}$$

In order to describe the mixing perturbatively we require $\rho < H$.

Quantization.—The fields are quantized in the interaction picture

$$\hat{\theta}_{\vec{k}}(\tau) = u_k(\tau)\hat{a}_{\vec{k}} + u_k^*(\tau)\hat{a}^\dagger_{-\vec{k}} \ , \tag{3.4}$$

$$\hat{\sigma}_{\vec{k}}(\tau) = v_k(\tau)\hat{b}_{\vec{k}} + v_k^*(\tau)\hat{b}^\dagger_{-\vec{k}} \ , \tag{3.5}$$

where $\mathrm{d}\tau = \mathrm{d}t/a(t)$ is conformal time and $u_k(\tau)$ and $v_k(\tau)$ are the free-field mode functions

$$u_k(\tau) = \frac{H}{\sqrt{2k^3}}(1 + ik\tau)e^{-ik\tau} \ , \tag{3.6}$$

$$v_k(\tau) = \sqrt{\frac{\pi}{2}}\frac{H}{\sqrt{2k^3}}(-k\tau)^{3/2}\, \mathrm{H}^{(1)}_\nu(-k\tau) \ , \qquad \text{with} \quad \nu \equiv \sqrt{\frac{9}{4} - \frac{m^2}{H^2}} \ . \tag{3.7}$$

Here, $\mathrm{H}^{(1)}_\nu$ is the Hankel function of the first kind. The massive mode decays on superhorizon scales, $v_k(\tau) \sim (-\tau)^{3/2-\nu}$. This feature completely explains the non-trivial scaling of correlation functions in the soft limits [25]. The mode functions have been normalized such that $[\hat{a}_{\vec{k}}, \hat{a}^\dagger_{\vec{k}'}] = [\hat{b}_{\vec{k}}, \hat{b}^\dagger_{\vec{k}'}] = (2\pi)^3\delta(\vec{k} + \vec{k}')$, which makes $\hat{a}_{\vec{k}}$ ($\hat{a}^\dagger_{\vec{k}}$) and $\hat{b}_{\vec{k}}$ ($\hat{b}^\dagger_{\vec{k}}$) the standard annihilation (creation) operators. Since there is no θ-σ coupling in $\mathcal{H}_0$, we also have $[\hat{a}_{\vec{k}}, \hat{b}^\dagger_{\vec{k}}] = 0$. We compute correlation functions of products of field operators, $Q(\tau)$, in the *in-in* formalism [55, 56]

$$\langle Q(\tau)\rangle = \langle\Omega|\left[\bar{\mathrm{T}}\exp\left(i\int_{-\infty}^{\tau}\mathrm{d}\tau'\,\hat{H}_{\rm int}(\tau')\right)\right]\hat{Q}(\tau)\left[\mathrm{T}\exp\left(-i\int_{-\infty}^{\tau}\mathrm{d}\tau'\,\hat{H}_{\rm int}(\tau')\right)\right]|\Omega\rangle \ , \tag{3.8}$$

where the symbols T and $\bar{\mathrm{T}}$ denote time-ordering and anti-time-ordering, respectively. The interaction Hamiltonian $H_{\rm int} = \int \mathrm{d}^3x\, a^4\, \mathcal{H}_{\rm int}$ can be written as a sum of two terms

$$H_2 \equiv -\int \mathrm{d}^3x\, a^3\rho\theta'\sigma \qquad \text{and} \qquad H_3 \equiv \int \mathrm{d}^3x\, a^4\mu\sigma^3 \ . \tag{3.9}$$

The mixing term H_2 is treated as a perturbative insertion in Feynman diagrams (see Fig. 5).



Figure 5: Perturbative description of N-point functions in quasi-single-field inflation.

Soft limits.—We end this review section with a brief summary of the soft limits of QSFI:

- The three-point function has the following momentum scaling in the squeezed limit
$$\lim_{k_1\to 0} \frac{\langle \zeta_{\vec{k}_1}\zeta_{\vec{k}_2}\zeta_{\vec{k}_3}\rangle'}{P_\zeta(k_1)} \ \propto\ k_1^{\frac{3}{2}-\nu}\ . \tag{3.10}$$
This result has a simple explanation [25]: The long (or soft) σ-mode exits the horizon at the time $\tau_1 \sim k_1^{-1}$, after which it decays with a rate that depends on its mass, $\sigma_k(\tau) \sim (-\tau)^{3/2-\nu}$. Its correlation with the short (or hard) modes is evaluated at the horizon crossing of the short modes, i.e. at $\tau_2 \sim k_2^{-1}$. By this time, the amplitude of the long σ-mode has decayed by a factor of $(\tau_2/\tau_1)^{3/2-\nu} = (k_1/k_2)^{3/2-\nu}$. This explains the suppression of the squeezed limit in eq. (3.10). Below we will confirm this intuition with an exact calculation.

- The four-point function in the collapsed limit scales as
$$\lim_{k_{12}\to 0} \frac{\langle \zeta_{\vec{k}_1}\zeta_{\vec{k}_2}\zeta_{\vec{k}_3}\zeta_{\vec{k}_4}\rangle'}{P_\zeta(k_{12})} \ \propto\ k_{12}^{3-2\nu}\ . \tag{3.11}$$
Again, the scaling follows from the superhorizon evolution of the soft σ-mode.

- The amplitudes of the collapsed four-point function and the squeezed three-point function are related as
$$\hat{\tau}_{\rm NL} \ \sim\ \frac{(\frac{6}{5}\hat{f}_{\rm NL})^2}{\left(\frac{\rho}{H}\right)^2} \ \gg\ (\tfrac{6}{5}\hat{f}_{\rm NL})^2\ . \tag{3.12}$$
We can understand this from the Feynman diagrams in Fig. 5. While in the three-point function every σ-propagator attaches to an external θ (or ζ), in the four-point function the virtual σ-mode does not have to be converted to ζ. The four-point function therefore only gets four powers of the small coupling ρ/H, while the square of the three-point function has six powers of ρ/H. For fixed three-point amplitude, we therefore get a boost in the four-point amplitude by a factor of $(\rho/H)^{-2} \gg 1$.

3.2 Explicit ‘in-in’ Computations

We now want to go beyond the qualitative discussion of the previous section, and compute the soft limits of the three- and four-point functions explicitly in terms of the fundamental parameters of the theory.

3.2.1 Power Spectrum

To assess the regime of validity of perturbation theory, we first use the *in-in* master formula (3.8) to compute the leading correction to the power spectrum of the adiabatic mode θ,

$$\langle \theta_{\vec{k}_1} \theta_{\vec{k}_2} \rangle = i^2 \int_{-\infty}^{0} \mathrm{d}\tau_1 \int_{-\infty}^{\tau_1} \mathrm{d}\tau_2 \, \langle [\hat{H}_2(\tau_2), [\hat{H}_2(\tau_1), \hat{\theta}_{\vec{k}_1} \hat{\theta}_{\vec{k}_2}]] \rangle \, . \tag{3.13}$$

Substituting eqs. (3.6), (3.7) and (3.9), we find [24]

$$\Delta_\zeta^2 \equiv k^3 P_\zeta(k) = \frac{1}{4} \frac{H^4}{M_{\rm pl}^2 |\dot{H}|} \left[1 + c(\nu) \left(\frac{\rho}{H} \right)^2 + \cdots \right] \, , \tag{3.14}$$

where

$$\begin{aligned} c(\nu) \equiv 2\pi \, \mathrm{Re} \Big[\int_0^\infty \mathrm{d}x_1 \int_{x_1}^\infty \mathrm{d}x_2 \, \Big(& x_1^{-1/2} \mathrm{H}_\nu^{(1)}(x_1) e^{i x_1} x_2^{-1/2} \mathrm{H}_\nu^{(2)}(x_2) e^{-i x_2} \\ & - x_1^{-1/2} \mathrm{H}_\nu^{(1)}(x_1) e^{-i x_1} x_2^{-1/2} \mathrm{H}_\nu^{(2)}(x_2) e^{-i x_2} \Big) \Big] \, , \end{aligned} \tag{3.15}$$

$=$ 

The function $c(\nu)$ formally diverges in the limit $\nu \to \frac{3}{2}$. In this limit, the isocurvaton σ becomes massless and freezes after horizon exit, instead of decaying. As a result, the transfer of σ to ζ continues for an indefinite time after horizon exit and evaluating the curvature perturbation at the infinite future $\tau = 0$ is no longer justified. Instead, the bispectrum should be evaluated at the time τ_f when inflation ends. This regulates the divergence.[17] Finally, we remark that the calculation is under perturbative control if and only if

$$c(\nu) \left(\frac{\rho}{H} \right)^2 < 1 \, . \tag{3.16}$$

3.2.2 Soft Limits: Methodology

We now derive the soft limits of QSFI in a style that will be familiar to cosmologists [4]. At the end of §3.2.3 and §3.2.4, we will demonstrate the precise equivalence between this method and the formalism of §2.

Long–short split.—In cosmology, soft limits are described as the correlation between long and short modes. Following [3, 4, 58], we treat the long-wavelength fields as *classical* backgrounds for the short-wavelength *quantum* fields. The fields θ and σ are split into long and short modes,[18]

$$\theta = \theta_L + \theta_S \quad \text{and} \quad \sigma = \sigma_L + \sigma_S, \tag{3.17}$$

---

[17] For the specific mixing term we consider, $\mathcal{H}_{\rm mix} = -\rho \dot{\theta} \sigma$, an effective mass of order ρ is generated for σ [24, 57]. If we were to re-sum the perturbation series, we expect that the divergence to be regulated by the effective mass. However, it isn't guaranteed that this possibility exists for generic mixing interactions, so we will ignore it here as well.

[18] This is morally similar to the split into soft and hard modes in the factorization theorems of QCD [59].

where

$$\theta_L(\vec{x},\tau) = \int_{k<\Lambda} \frac{\mathrm{d}^3k}{(2\pi)^3}\, \theta_{\vec{k}}(\tau) e^{i\vec{k}.\vec{x}} \qquad \text{and} \qquad \theta_S(\vec{x},\tau) = \int_{k>\Lambda} \frac{\mathrm{d}^3k}{(2\pi)^3}\, \theta_{\vec{k}}(\tau) e^{i\vec{k}.\vec{x}} \ , \tag{3.18}$$

and similarly for σ. The precise value of the momentum cut-off Λ won't be important. We then write the interaction Hamiltonian $H_{\rm int}$ in terms of θ_L, θ_S, σ_L and σ_S. The quadratic mixing term H_2 doesn't produce any coupling between long and short modes. The only relevance of the mixing is therefore to convert σ_S into θ_S via

$$H_{SS} = -\rho a^3 \int \mathrm{d}^3x\, \sigma_S \theta_S' \ . \tag{3.19}$$

The cubic interaction H_3, on the other hand, does couple long modes to short modes. The only interaction that will be relevant in the soft limit is

$$H_{LSS} = 3\mu a^4 \int \mathrm{d}^3x\, \sigma_L \sigma_S^2 \ . \tag{3.20}$$

Soft limits.—Our goal is to use eqs. (3.19) and (3.20) to compute the squeezed limit of the three-point function,

$$\langle \theta_{\vec{k}_1} \theta_{\vec{k}_2} \theta_{\vec{k}_3} \rangle \xrightarrow{k_1 \to 0} \langle \theta_{\vec{k}_1} \langle \theta_{\vec{k}_2} \theta_{\vec{k}_3} \rangle_{\sigma_L} \rangle \ , \tag{3.21}$$

and the collapsed limit of the four-point function,

$$\langle \theta_{\vec{k}_1} \theta_{\vec{k}_2} \theta_{\vec{k}_3} \theta_{\vec{k}_4} \rangle \xrightarrow{k_{12} \to 0} \langle \langle \theta_{\vec{k}_1} \theta_{\vec{k}_2} \rangle_{\sigma_L} \langle \theta_{\vec{k}_3} \theta_{\vec{k}_4} \rangle_{\sigma_L} \rangle \ . \tag{3.22}$$

We see that both soft limits can be expressed in terms of the modulated power spectrum

$$\langle \theta_{\vec{k}_1} \theta_{\vec{k}_2} \rangle_{\sigma_L} \ , \tag{3.23}$$

i.e. the power spectrum $\langle \theta_{\vec{k}_1} \theta_{\vec{k}_2} \rangle$ in the presence of a long-wavelength background σ_L. We compute eq. (3.23) in the *in-in* formalism

$$\langle \theta_{\vec{k}_1} \theta_{\vec{k}_2} \rangle_{\sigma_L(\tau)} = i^3 \int_{-\infty}^{\tau} \mathrm{d}\tau_1 \int_{-\infty}^{\tau_1} \mathrm{d}\tau_2 \int_{-\infty}^{\tau_2} \mathrm{d}\tau_3\ \langle \Big[\hat{H}_{\rm int}(\tau_3), \Big[\hat{H}_{\rm int}(\tau_2), \Big[\hat{H}_{\rm int}(\tau_1), \hat{\theta}_{\vec{k}_1} \hat{\theta}_{\vec{k}_2}(\tau)\Big]\Big]\Big] \rangle \ , \tag{3.24}$$

where $H_{\rm int} = H_{LSS} + H_{SS}$. In the limit $\tau \to 0$, this becomes

$$\langle \theta_{\vec{k}_1} \theta_{\vec{k}_2} \rangle_{\sigma_L} = -\frac{3\pi^2}{2} \frac{\mu \rho^2}{H^2} \frac{1}{k_1^3} \left(A_{12} + B_{12} \right) \ , \tag{3.25}$$

where the A-term is obtained by replacing $H_{\rm int}(\tau_3)$ by H_{LSS} and the two other interaction terms by H_{SS}, while the B-term is obtained by replacing $H_{\rm int}(\tau_2)$ by H_{LSS} and the rest by H_{SS}. This leads to

$$A_{12} \equiv \int_{-\infty}^{0} \mathrm{d}x_1 \int_{-\infty}^{x_1} \mathrm{d}x_2 \int_{-\infty}^{x_2} \mathrm{d}x_3\ f_A(x_1,x_2,x_3)\ (\sigma_L)_{\vec{k}_1+\vec{k}_2}(x_3/k_1) \ , \tag{3.26}$$

$$B_{12} \equiv \int_{-\infty}^{0} \mathrm{d}x_1 \int_{-\infty}^{x_1} \mathrm{d}x_2 \int_{-\infty}^{x_2} \mathrm{d}x_3\ f_B(x_1,x_2,x_3)\ (\sigma_L)_{\vec{k}_1+\vec{k}_2}(x_2/k_1) \ , \tag{3.27}$$

with

$$f_A(x_i) \equiv (-x_1 x_2 (x_3)^2)^{-1/2} \sin(-x_1) \sin(-x_2) \times\ \mathrm{Im}\Big(\mathrm{H}_\nu^{(2)}(-x_1)\, \mathrm{H}_\nu^{(2)}(-x_2) (\mathrm{H}_\nu^{(1)}(-x_3))^2\Big) \ ,$$

$$f_B(x_i) \equiv (-x_1 (x_2)^2 x_3)^{-1/2} \sin(-x_1) \times\ \mathrm{Im}\Big(\mathrm{H}_\nu^{(1)}(-x_1)\, \mathrm{H}_\nu^{(2)}(-x_2)\Big) \mathrm{Im}\Big(\mathrm{H}_\nu^{(1)}(-x_2)\, \mathrm{H}_\nu^{(2)}(-x_3) e^{ix_3}\Big) \ .$$

3.2.3 Bispectrum in the Squeezed Limit

With this preliminary work, it is now straightforward to compute the squeezed limit of the bispectrum,

$$\langle \theta_{\vec{k}_1} \langle \theta_{\vec{k}_2} \theta_{\vec{k}_3} \rangle_{\sigma_L} \rangle = -\frac{3\pi^2}{2} \frac{\mu\rho^2}{H^2} \frac{1}{k_2^3} \Big(\langle \theta_{\vec{k}_1} A_{23} \rangle + \langle \theta_{\vec{k}_1} B_{23} \rangle \Big) \, . \tag{3.28}$$

The only slight complication is due to the fact that the calculation requires the unequal time correlation function of two fields, i.e. $\langle \sigma_{\vec{k}}(\tau) \theta_{\vec{q}}(0) \rangle$. However, since $\theta_{\vec{q}}(\tau)$ freezes in the limit $|q\tau| \ll 1$, we have $\lim\limits_{|q\tau| \ll 1} \theta_{\vec{q}}(\tau) \approx \theta_{\vec{q}}(0)$ and hence

$$\langle \sigma_{\vec{k}}(\tau) \theta_{\vec{q}}(0) \rangle \xrightarrow{|k\tau| \ll 1} \langle \sigma_{\vec{k}}(\tau) \theta_{\vec{q}}(\tau) \rangle \, , \tag{3.29}$$

where

$$\begin{aligned} \langle \sigma_{\vec{k}}(\tau) \theta_{\vec{q}}(\tau) \rangle &= -i \int_{-\infty}^{\tau} \mathrm{d}\tilde{\tau} \int_{\vec{p}} a(\tilde{\tau})^3 \rho \, \langle 0 | \hat{\sigma}_{\vec{p}}(\tilde{\tau}) \hat{\theta}'_{-\vec{p}}(\tilde{\tau}) \hat{\sigma}_{\vec{k}}(\tau) \hat{\theta}_{\vec{q}}(\tau) | 0 \rangle \; + \; c.c. \\ &= -2\rho \, (2\pi)^3 \delta(\vec{k} + \vec{q}) \, \mathrm{Re} \left[i \int_{-\infty}^{\tau} \mathrm{d}\tilde{\tau} \, a(\tilde{\tau})^3 \, v_k(\tilde{\tau}) v_k^*(\tau) u_q'(\tilde{\tau}) u_q^*(\tau) \right] \, . \end{aligned} \tag{3.30}$$

Using the superhorizon limits of the mode functions,

$$\lim_{|k\tau| \ll 1} u_k(\tau) \approx \frac{H}{\sqrt{2k^3}} \qquad \text{and} \qquad \lim_{|k\tau| \ll 1} v_k(\tau) \approx -i \, \frac{2^{\nu} \Gamma(\nu)}{2\sqrt{\pi}} \frac{H}{\sqrt{k^3}} (-k\tau)^{3/2 - \nu} \, , \tag{3.31}$$

we get

$$\langle \sigma_{\vec{k}}(\tau) \theta_{\vec{q}}(\tau) \rangle \xrightarrow{|k\tau| \ll 1} \sqrt{\frac{2}{\pi}} \frac{\Gamma(\nu)}{2^{2-\nu}} \frac{\rho}{H} \frac{H^2}{k^3} (-k\tau)^{3/2 - \nu} \, d(\nu) \, (2\pi)^3 \delta(\vec{p} + \vec{q}) \, , \tag{3.32}$$

where

$$d(\nu) \equiv \sqrt{\frac{\pi}{2}} \int_{-\infty}^{0} \mathrm{d}x \, (-x)^{-1/2} \, \mathrm{Re} \left[\mathrm{H}_{\nu}^{(1)}(-x) e^{-ix} \right] \; = \tag{3.33}$$



Substituting (3.32) into (3.28), we obtain the squeezed limit of the bispectrum for the curvature perturbation ζ,

$$\langle \zeta_{\vec{k}_1} \zeta_{\vec{k}_2} \zeta_{\vec{k}_3} \rangle' \xrightarrow{k_1 \to 0} \frac{12}{5} f_{\mathrm{NL}}(\nu) \cdot P_{\zeta}(k_1) P_{\zeta}(k_2) \left(\frac{k_1}{k_2} \right)^{3/2 - \nu} \, , \tag{3.34}$$

where

$$f_{\mathrm{NL}}(\nu) \equiv p(\nu) \, d(\nu) \cdot \frac{\mu\rho^3}{H^4} \cdot \Delta_{\zeta}^{-1} \, . \tag{3.35}$$

Here, $d(\nu)$ is the same function as in eq. (3.33) and $p(\nu)$ is defined as

$$p(\nu) \equiv 5\pi^{3/2}\,\frac{\Gamma(\nu)}{2^{3-\nu}}\int_{-\infty}^{0}{\rm d}x_1\int_{-\infty}^{x_1}{\rm d}x_2\int_{-\infty}^{x_2}{\rm d}x_3\,\Big[(-x_3)^{3/2-\nu}f_A(x_i)+(-x_2)^{3/2-\nu}f_B(x_i)\Big]\ . \tag{3.36}$$

Eq. (3.34) precisely matches the result that one obtains by taking the limit $k_1 \to 0$ of the full expression of the bispectrum [24].

In the following insert we show how to reproduce eq. (3.34) from our analysis in §2:

We start by defining a complete set of one-particle states

$$|1_{\vec q}^{(\zeta)}\rangle \equiv P_\zeta^{-1/2}|\zeta_{\vec q}\rangle \quad \text{and} \quad |1_{\vec q}^{(\tilde\sigma)}\rangle \equiv P_{\tilde\sigma}^{-1/2}|\tilde\sigma_{\vec q}\rangle\ . \tag{3.37}$$

In order for the states to be orthogonal, we have defined

$$\tilde\sigma_{\vec q} \equiv \sigma_{\vec q} - \frac{\langle\zeta_{\vec q}|\sigma_{\vec q}\rangle'}{P_\zeta(q)}\,\zeta_{\vec q}\ . \tag{3.38}$$

As our expansion parameter we define

$$\varepsilon^2(\nu) \equiv \frac{|\langle\zeta\sigma\rangle'|^2}{P_\zeta P_\sigma} < 1\ . \tag{3.39}$$

To determine the effect of σ on the squeezed limit of the three-point function of ζ, we first compute the parameter $c_{\tilde\sigma}$ in eq. (2.40). Since spatial dilations act linearly on σ, we have

$$i\langle[Q_d,\tilde\sigma_{\vec q}]\rangle = \frac{\langle\zeta_{\vec q}|\sigma_{\vec q}\rangle'}{P_\zeta(q)}(2\pi)^3\delta(\vec q)\ . \tag{3.40}$$

This implies that

$$c_{\tilde\sigma}(q) = -i\frac{\langle\zeta_{\vec q}|\sigma_{\vec q}\rangle'}{P_\zeta P_{\tilde\sigma}^{1/2}}\ . \tag{3.41}$$

The coefficient $c_{\tilde\sigma}$ may have a non-zero real part, but it won't contribute in what follows. We therefore get

$${\rm Im}\Big[c_{\tilde\sigma}(q)\langle 1_{\vec q}^{(\tilde\sigma)}|\zeta^2\rangle'\Big] \ = \ -\frac{\langle\zeta_{\vec q}|\sigma_{\vec q}\rangle'}{P_\zeta P_{\tilde\sigma}}\,{\rm Im}\big[i\langle\tilde\sigma_{\vec q}|\zeta^2\rangle'\big] \ \simeq \ -\frac{\langle\zeta_{\vec q}|\sigma_{\vec q}\rangle'\langle\sigma_{\vec q}|\zeta^2\rangle'}{P_\zeta P_\sigma}\big[1+\mathcal{O}(\varepsilon^2)\big]\ , \tag{3.42}$$

and eq. (2.40) becomes

$$\frac{\langle\zeta_{\vec q}\zeta_{\vec k_1}\zeta_{\vec k_2}\rangle'}{P_\zeta(q)} \xrightarrow{q\to 0} \frac{\langle\zeta_{\vec k}|\sigma_{\vec q}\rangle'\langle\sigma_{\vec q}|\zeta_{\vec k_1}\zeta_{\vec k_2}\rangle'}{P_\zeta(q)P_\sigma(q)}\ . \tag{3.43}$$

This is equivalent to the starting point of this section, eq. (3.28), so following the same steps will lead to the answer in eq. (3.34).

3.2.4 Trispectrum in the Collapsed Limit

The collapsed limit of the four-point function is equal to the correlation of two modulated power spectra,

$$\langle \langle \theta_{\vec{k}_1} \theta_{\vec{k}_2} \rangle_{\sigma_L} \langle \theta_{\vec{k}_3} \theta_{\vec{k}_4} \rangle_{\sigma_L} \rangle = \left(\frac{3\pi^2}{2} \frac{\mu \rho^2}{H^2} \right)^2 \frac{1}{k_1^3 k_3^3} \Big(\underbrace{\langle A_{12} A_{34} \rangle + \langle A_{12} B_{34} \rangle + \langle B_{12} A_{34} \rangle + \langle A_{12} B_{34} \rangle}_{(\star)} \Big) \ , \tag{3.44}$$

where the sum over contractions is

$$\begin{aligned} (\star) \ = \ & \left(\prod_{i=1}^{3} \int \mathrm{d}x_i \right) \left(\prod_{i=1}^{3} \int \mathrm{d}y_i \right) \times \\ & \times \Big[\ f_A(x_i) f_A(y_i) \left\langle (\sigma_L)_{\vec{k}_{12}}(\tfrac{x_3}{k_1}) \, (\sigma_L)_{\vec{k}_{34}}(\tfrac{y_3}{k_3}) \right\rangle \ + \ f_A(x_i) f_B(y_i) \left\langle (\sigma_L)_{\vec{k}_{12}}(\tfrac{x_3}{k_1}) \, (\sigma_L)_{\vec{k}_{34}}(\tfrac{y_2}{k_3}) \right\rangle \\ & \ + \ f_B(x_i) f_A(y_i) \left\langle (\sigma_L)_{\vec{k}_{12}}(\tfrac{x_2}{k_1}) \, (\sigma_L)_{\vec{k}_{34}}(\tfrac{y_3}{k_3}) \right\rangle \ + \ f_B(x_i) f_B(y_i) \left\langle (\sigma_L)_{\vec{k}_{12}}(\tfrac{x_2}{k_1}) \, (\sigma_L)_{\vec{k}_{34}}(\tfrac{y_2}{k_3}) \right\rangle \ \Big] \ . \end{aligned} \tag{3.45}$$

Using

$$\langle \sigma_{\vec{q}_1}(\tau_1) \sigma_{\vec{q}_2}(\tau_2) \rangle \ \xrightarrow{q_1, q_2 \to 0} \ \frac{2^{2\nu-1}\Gamma(\nu)^2}{\pi} \frac{H^2}{2 q_1^3} (-q_1 \tau_1)^{3/2-\nu} (-q_2 \tau_2)^{3/2-\nu} \ (2\pi)^3 \delta\left(\vec{q}_1 + \vec{q}_2\right) \ , \tag{3.46}$$

this becomes

$$\begin{aligned} (\star) \ \to \ & \frac{\Gamma(\nu)^2}{2^{2-2\nu}\pi} \frac{H^2}{(k_1 k_3)^{3/2-\nu}} \frac{1}{(k_{12})^{2\nu}} \, (2\pi)^3 \, \delta\big(\vec{k}_1 + \vec{k}_2 + \vec{k}_3 + \vec{k}_4\big) \times \\ & \times \left[\left(\prod_{i=1}^{3} \int \mathrm{d}x_i \right) \Big(f_A(x_i) \, (-x_3)^{3/2-\nu} + f_B(x_i) \, (-x_2)^{3/2-\nu} \Big) \right]^2 \ . \end{aligned} \tag{3.47}$$

Eq. (3.44) can therefore be written as

$$\langle \theta_{\vec{k}_1} \theta_{\vec{k}_2} \theta_{\vec{k}_3} \theta_{\vec{k}_4} \rangle' \ \xrightarrow{k_{12} \to 0} \ t(\nu) \cdot \frac{\rho^4 \mu^2}{H^6} \frac{H^4}{(k_1 k_3)^{9/2-\nu}} \frac{1}{(k_{12})^{2\nu}} \ , \tag{3.48}$$

where

$$t(\nu) \equiv \left(\tfrac{6}{5} \, p(\nu) \right)^2 \ = \tag{3.49}$$



The function $t(\nu)$ diverges as $\nu \to 0$ and $\nu \to \frac{3}{2}$. The divergence for $\nu \to 0$ arises from the fact that the expansion of the Hankel function in eq. (3.31) isn't justified when $k_{12}/k \ll e^{-1/\nu}$. Instead the expansion of the Hankel function should be changed to

$$v_{k_{12}}(x/k_1) \to -i \sqrt{\frac{\pi}{2}} \frac{H}{\sqrt{2 k_{12}^3}} \left(-\frac{k_{12}}{k_1} x \right)^{3/2} \ln \left(\frac{k_{12}}{k_1} \right) \ . \tag{3.50}$$

This regulates the divergence. The divergence of $t(\nu)$ for $\nu \to \frac{3}{2}$ has the same origin (and the same resolution) as the divergence of $c(\nu)$ in (3.15). Finally, converting (3.48) to curvature perturbations, we get

$$\boxed{\langle\zeta_{\vec{k}_1}\zeta_{\vec{k}_2}\zeta_{\vec{k}_3}\zeta_{\vec{k}_4}\rangle' \xrightarrow{k_{12}\to 0} 4\,\tau_{\rm NL}(\nu)\cdot P_\zeta(k_1)P_\zeta(k_3)P_\zeta(k_{12})\left(\frac{k_{12}}{(k_1k_3)^{1/2}}\right)^{3-2\nu}}\ , \tag{3.51}$$

where

$$\boxed{\tau_{\rm NL}(\nu) = \frac{(\frac{6}{5}f_{\rm NL}(\nu))^2}{\varepsilon^2(\nu)}}\ , \qquad \text{with} \qquad \varepsilon(\nu) \equiv \frac{\rho}{H}d(\nu)\ . \tag{3.52}$$

Note that we have recovered the scaling in eq. (3.11). Since $d^2(\nu) \simeq c(\nu)$, the condition on perturbative control, eq. (3.16), becomes

$$\varepsilon(\nu) = \frac{\rho}{H}d(\nu) < 1\ . \tag{3.53}$$

Hence, as long as the calculation is under perturbative control, we find $\tau_{\rm NL} \geq (\frac{6}{5}f_{\rm NL})^2$, consistent with the Suyama-Yamaguchi inequality [15].

Finally, we show how eq. (3.51) also follows from our analysis in §2:

We insert the complete set of one-particle states, eq. (3.37), into the four-point function

$$\langle\zeta^4\rangle = \int_{\vec{q}} \Big[\langle\zeta^2|1_{\vec{q}}^{(\zeta)}\rangle\langle 1_{\vec{q}}^{(\zeta)}|\zeta^2\rangle + \langle\zeta^2|1_{\vec{q}}^{(\tilde{\sigma})}\rangle\langle 1_{\vec{q}}^{(\tilde{\sigma})}|\zeta^2\rangle\Big]\ , \tag{3.54}$$

where

$$|1_{\vec{q}}^{(\tilde{\sigma})}\rangle = N\Big[|1_{\vec{q}}^{(\sigma)}\rangle - \alpha\,|1_{\vec{q}}^{(\zeta)}\rangle\Big]\ , \tag{3.55}$$

with $\alpha = \langle 1_{\vec{q}}^{(\zeta)}|1_{\vec{q}}^{(\sigma)}\rangle' \sim \varepsilon(\nu)$ and $N = 1 + \mathcal{O}(\varepsilon)$. At leading order in ε, eq. (3.54) becomes

$$\langle\zeta_{\vec{k}_1}\zeta_{\vec{k}_2}\zeta_{\vec{k}_3}\zeta_{\vec{k}_4}\rangle' \xrightarrow{k_{12}\to 0} \frac{1}{P_\sigma(k_{12})}\langle\zeta_{\vec{k}_1}\zeta_{\vec{k}_2}\sigma_{-\vec{k}_{12}}\rangle'\langle\sigma_{-\vec{k}_{34}}\zeta_{\vec{k}_3}\zeta_{\vec{k}_4}\rangle'\ . \tag{3.56}$$

This is equivalent to the starting point of this section, eq. (3.44), so following the same steps will lead to the answer in eq. (3.51).

---

4 Discussion: Non-Gaussianity as a Particle Detector



$\lim_{k\to 0} k^3\langle\zeta_{\vec{k}}\zeta_{\vec{k}_1}\zeta_{\vec{k}_2}\rangle$	k^2	$k^{\frac{3}{2}-\nu}$	k^0
$\hat{\tau}_{\rm NL}$	$= \left(\frac{6}{5}\hat{f}_{\rm NL}\right)^2$	$\gg \left(\frac{6}{5}\hat{f}_{\rm NL}\right)^2$	$\geq \left(\frac{6}{5}\hat{f}_{\rm NL}\right)^2$

Figure 6: Particle spectra of inflationary models and their observational signatures.

Observations of the CMB and LSS probe the physics of inflation through the statistics of the primordial curvature perturbations. Soft limits of the correlation functions of these perturbations are especially sensitive to the number of light degrees of freedom during inflation. In this paper, we have explained how the spectrum of particles can impact observations (see Fig. 6).

Single-field inflation.—The fluctuations in single-field inflation are Goldstone bosons of a spontaneously broken dilation symmetry [11]. Being non-linearly realized, the broken symmetry is still reflected in Ward identities relating the squeezed limits of N-point functions to the symmetry variation of $(N-1)$-point functions [11, 42, 43]. Up to small corrections associated with the breaking of scale-invariance, the squeezed limits of all N-point functions therefore vanish in single-field inflation [3]. This is analogous to the Adler zero [26] of chiral perturbation theory [29]. In this note, we have given a new proof of this famous result. Measuring a significant three-point function in the squeezed limit would therefore automatically rule out *all* single-field models. Going beyond single-field inflation, the squeezed limit captures information about the spectrum of additional light degrees of freedom.

Multi-field inflation.—Coupling additional fields to the inflaton relaxes some of the symmetry constraints on inflationary correlation functions. The extra fields can therefore produce a significant signal in the squeezed limit. In fact, if the non-Gaussianity arises from superhorizon evolution, then the signal tends to be peaked in the squeezed configuration. If a single field (but not necessarily the inflaton) creates the primordial curvature perturbations ζ and their non-Gaussianity, then the collapsed limit of the four-point function is uniquely fixed by the squeezed limit of the three-point function. However, this correlation is broken if more than one field sources ζ and its interactions. In that case one can show that the collapsed four-point function is always larger than the square of the squeezed three-point function [15]. In this note, we have given a new proof of this result.

Quasi-single-field inflation.—Supersymmetric models of inflation often mix a light inflaton field with an additional isocurvaton field whose mass is of order the inflationary Hubble scale [25]. Such models of quasi-single-field inflation [24] naturally give rise to a boosted four-point function and to characteristic momentum scalings in the soft limits. We have computed these signatures explicitly in terms of the fundamental parameters of the theory. As we have explained elsewhere [22, 25], these results are relevant for connecting the microscopic physics of QSFI to late-time observables such as stochastic halo bias [22].

The above shows that soft limits of primordial non-Gaussianity provide a remarkable opportunity to probe the spectrum of particles during inflation, i.e. at energies totally inaccessible to conventional particle physics experiments. The PLANCK satellite [60] will measure the CMB anisotropies over a large range of scales and hence will significantly improve the observational constraints on the soft limits of primordial non-Gaussianity. At the same time, the Sloan Digital Sky Survey [61] continues to collect data on the clustering of collapsed objects. The scale-dependent biasing [1] of these objects is a very sensitive probe of the squeezed limit of the primordial correlation functions [62]. Looking ahead, LSS-experiments like EUCLID [63] and LSST [64], as well as CMB-experiments like PIXIE [65] and CMBPol [66] all have the potential to teach us a great deal about the soft limits of inflationary correlations.

Acknowledgments

Xingang Chen, Paolo Creminelli, Justin Khoury, Marko Simonović and Gianmassimo Tasinato kindly provided comments on a draft of the paper. We thank Raphael Flauger and Rafael Porto for helpful discussions. D.B. and V.A. gratefully acknowledge support from a Starting Grant of the European Research Council (ERC STG grant 279617). The research of D.G. is supported by the DOE under grant number DE-FG02-90ER40542 and the Martin A. and Helen Chooljian Membership at the Institute for Advanced Study. D.B. thanks the theory groups at Groningen, Edinburgh, Oxford, IUCAA (Pune), TIFR (Mumbai) and UCL (London) for their kind hospitality and the opportunity to present this work.

A Consistency Relations at Finite Momentum

The soft limits described in §2.2 were presented in the strict $q \to 0$ limit. However, since these results follow from current conservation alone, one might expect that they can be extended to finite q. In this appendix, we will demonstrate that this is indeed the case.

By definition, any conserved charge is written as an integral of the current over a spatial slice. Throughout the main text, we have worked directly with the charge $Q \equiv \int \mathrm{d}^3x\, J^0(\vec{x},t)$ to derive our results. In particular, we have considered the following commutator

$$i\langle[Q,\zeta(\vec{y})\zeta(\vec{z})]\rangle \;=\; \langle\delta_Q\zeta(\vec{y})\zeta(\vec{z})\rangle + \langle\zeta(\vec{y})\delta_Q\zeta(\vec{z})\rangle \;, \tag{A.1}$$

where it is understood that all fields for which the time dependence isn't shown are evaluated at the same time $t_\star$. Eq. (A.1) follows from current conservation via the Ward identity [30, 50]

$$\begin{aligned} i\,\partial_\mu^{(x)}\langle J^\mu(\vec{x},t)\zeta(\vec{y},t_\star)\zeta(\vec{z},t_\star,)\rangle_T \;&=\; \delta_{\mathcal{C}}(t-t_\star)\delta(\vec{x}-\vec{y})\,\langle\delta_Q\zeta(\vec{y},t_\star)\zeta(\vec{z},t_\star)\rangle \\ &\quad+\,\delta_{\mathcal{C}}(t-t_\star)\delta(\vec{x}-\vec{z})\,\langle\zeta(\vec{y},t_\star)\delta_Q\zeta(\vec{z},t_\star)\rangle \;, \end{aligned} \tag{A.2}$$

where the terms proportional to $\delta_{\mathcal{C}}(t-t_\star)$ arise from the time derivative acting on the "time-ordering" operator T. This feature of the Ward identity deserves a few more words of explanation: By definition, the above *in-in* correlation functions are "time-ordered" on a contour $\mathcal{C}$ that runs from $t = -\infty + i\epsilon$ to $-\infty - i\epsilon$, passing through $t_\star$. Formally, we can write this as

$$\langle\mathcal{O}_1(t_1)\mathcal{O}_2(t_2)\rangle_T \;=\; \theta(\lambda_1-\lambda_2)\,\langle\mathcal{O}_1(t_1)\mathcal{O}_2(t_2)\rangle + \theta(\lambda_2-\lambda_1)\,\langle\mathcal{O}_2(t_2)\mathcal{O}_1(t_1)\rangle \;, \tag{A.3}$$

where θ is the Heaviside function and λ is a real number that parametrizes the contour $\mathcal{C}$—e.g. we may choose $\lambda = t - t_\star$ on the "incoming" contour $\mathcal{C}_+$ and $\lambda = t_\star - t$ on the "outgoing" contour $\mathcal{C}_-$ (see Fig. 7). The delta function $\delta_{\mathcal{C}}(t-t_\star)$ on the r.h.s. of (A.2) is therefore defined as

$$\delta_{\mathcal{C}}(t-t_\star) = \begin{cases} \;\;\delta(t-t_\star) & \text{on } \mathcal{C}_+\,, \\ -\delta(t_\star-t) & \text{on } \mathcal{C}_-\,. \end{cases} \tag{A.4}$$

If we now integrate (A.2) over the contour $\int_{t_\star-\Delta t+i\epsilon}^{t_\star-\Delta t-i\epsilon}\mathrm{d}t\int\mathrm{d}^3x \equiv \oint\mathrm{d}t\int\mathrm{d}^3x$ through $t = t_\star$, we arrive at eq. (A.1). We have dropped the integral over $\partial_i J^i$ because it only receives contributions from spatial infinity.



Figure 7: Illustration of the contour employed in the computation of *in-in* correlation functions.

When translated into momentum space, the above treatment is limited to $q = 0$. However, nothing required that we integrate the Ward identity in this way. Alternatively, we can multiply eq. (A.2) by $e^{i\vec{q}\cdot\vec{x}}$ and then integrate over $\vec{x}$ and t,

$$\oint \mathrm{d}t \int \mathrm{d}^3x \; e^{i\vec{q}\cdot\vec{x}} \; i\,\partial_\mu^{(x)}\langle J^\mu(\vec{x},t)\zeta(\vec{y})\zeta(\vec{z})\rangle_T \;=\; e^{i\vec{q}\cdot\vec{y}}\langle\delta_Q\zeta(\vec{y})\zeta(\vec{z})\rangle + e^{i\vec{q}\cdot\vec{z}}\langle\zeta(\vec{y})\,\delta_Q\zeta(\vec{z})\rangle \;. \tag{A.5}$$

Taking the limit $\epsilon, \Delta t \to 0$ and going to momentum space, we find

$$i\langle[J^0_{\vec{q}}(t_\star), \zeta_{\vec{k}_1}\zeta_{\vec{k}_2}]\rangle + \oint \mathrm{d}t \; q_i\langle J^i_{\vec{q}}(t)\zeta_{\vec{k}_1}\zeta_{\vec{k}_2}\rangle = \langle\delta_Q\zeta_{\vec{q}+\vec{k}_1}\,\zeta_{\vec{k}_2}\rangle + \langle\zeta_{\vec{k}_1}\,\delta_Q\zeta_{\vec{q}+\vec{k}_2}\rangle \;. \tag{A.6}$$

Inserting a complete set of states in the first term, one arrives at the identity

$$\sum_{n,\vec{p}} i\Big[\langle J^0_{\vec{q}}(t_\star)|n_{\vec{p}}\rangle\langle n_{\vec{p}}|\zeta_{\vec{k}_1}\zeta_{\vec{k}_2}\rangle - \text{h.c.}\Big] \;=\; \langle\delta_Q\zeta_{\vec{q}+\vec{k}_1}\,\zeta_{\vec{k}_2}\rangle + \langle\zeta_{\vec{k}_1}\,\delta_Q\zeta_{\vec{q}+\vec{k}_2}\rangle - \oint \mathrm{d}t\; q_i\langle J^i_{\vec{q}}(t)\zeta_{\vec{k}_1}\zeta_{\vec{k}_2}\rangle \;. \tag{A.7}$$

Note that this result holds without making any assumptions about the relative or absolute sizes of q, k_1 and k_2. This isn't surprising, since so far we have only used current conservation which holds at all momenta. In order to arrive back at eq. (A.1), we must show that we can indeed neglect the extra term in eq. (A.7) in the limit $q \to 0$. The choice of contour in the complex time domain implies that

$$\oint \mathrm{d}t\; q_i\langle J^i_{\vec{q}}(t)\zeta_{\vec{k}_1}\zeta_{\vec{k}_2}\rangle \;\xrightarrow{\epsilon\to 0}\; \int_{t_\star-\Delta t}^{t_\star} \mathrm{d}t\; q_i\langle[J^i_{\vec{q}}(t), \zeta_{\vec{k}_1}\zeta_{\vec{k}_2}]\rangle \;. \tag{A.8}$$

Causality requires that $[J^i(\vec{x},t_\star), \zeta(\vec{y},t_\star)\zeta(\vec{z},t_\star)]$ vanishes when $\vec{x}$ is outside the light-cone[19] of $\vec{y}$ and $\vec{z}$. For $\vec{x} \neq \vec{y}, \vec{z}$, the operators are space-like separated and the commutator is a proportional to a delta function in position space, or in momentum space $[J^i_{\vec{q}}, \zeta_{\vec{k}}] = f(q)\mathcal{O}_{\vec{k}+\vec{q}}$, where $f(q)$ is an analytic function. Translation invariance ensures that the correlation function $\langle\mathcal{O}_{\vec{k}_1+\vec{q}}\,\zeta_{\vec{k}_2}\rangle$ can be written in terms of $\vec{k}_2$. Therefore, the final term in (A.8) will be analytic in q. Furthermore, invariance under spatial rotations requires that the commutator vanishes at $\vec{q} = 0$ (using the rotation $\vec{k}_1 \to -\vec{k}_1$). These two facts together imply

$$\lim_{q\to 0}\,\langle[J^i_{\vec{q}}(t), \zeta_{\vec{k}_1}\zeta_{\vec{k}_2}]\rangle = f(k_1,k_2)\,q^i \;. \tag{A.9}$$

Hence, we find

$$\lim_{q\to 0}\,\sum_{n,\vec{p}} i\Big[\langle J^0_{\vec{q}}|n_{\vec{p}}\rangle\langle n_{\vec{p}}|\zeta_{\vec{k}_1}\zeta_{\vec{k}_2}\rangle - \text{h.c.}\Big] \;=\; \langle\delta_Q\zeta_{\vec{q}+\vec{k}_1}\,\zeta_{\vec{k}_2}\rangle + \langle\zeta_{\vec{k}_1}\,\delta_Q\zeta_{\vec{q}+\vec{k}_2}\rangle + \mathcal{O}(q^2) \;. \tag{A.10}$$

The conclusion that the consistency relation holds at finite q up to corrections suppressed by q^2 is extremely important. It ensures that contributions to the l.h.s. of (A.10) that vanish like $q^{3-\alpha}$ for $\alpha > 1$ cannot be mimicked by finite q effects in single-field inflation. See [6] for an alternative derivation of this result.

---

[19] The causal structure of the background will be modified by metric fluctuations. In all situations of interest in this paper, the fluctuations are small and this effect is negligible [34]. On the other hand, in eternal inflation the presence of large metric fluctuations can modify the causal structure significantly [67].

References

[1] N. Dalal, O. Dore, D. Huterer, and A. Shirokov, "The Imprints of Primordial Non-Gaussianities on Large-Scale Structure," Phys. Rev. D **77**, 123514 (2008).

[2] E. Pajer and M. Zaldarriaga, "A New Window on Primordial Non-Gaussianity," arXiv:1201.5375 [astro-ph.CO].

[3] J. Maldacena, "Non-Gaussian Features of Primordial Fluctuations in Single-Field Inflationary Models," JHEP **0305**, 013 (2003).

[4] P. Creminelli and M. Zaldarriaga, "Single-Field Consistency Relation for the Three-Point Function," JCAP **0410**, 006 (2004).

[5] C. Cheung, A. Fitzpatrick, J. Kaplan, and L. Senatore, "On the Consistency Relation of the Three-Point Function in Single-Field Inflation," JCAP **0802**, 021 (2008).

[6] P. Creminelli, G. D'Amico, M. Musso, and J. Noreña, "The (not so) Squeezed Limit of the Primordial Three-Point Function," JCAP **1111**, 038 (2011).

[7] X. Chen, M. Huang, and G. Shiu, "The Inflationary Trispectrum for Models with Large Non-Gaussianities," Phys. Rev. D **74**, 121301 (2006).

[8] M. Li and Y. Wang, "Consistency Relations for Non-Gaussianity," JCAP **0809**, 018 (2008).

[9] L. Leblond and E. Pajer, "Resonant Trispectrum and a Dozen More Primordial N-Point Functions," JCAP **1101**, 035 (2011).

[10] D. Seery, M. Sloth, and F. Vernizzi, "Inflationary Trispectrum from Graviton Exchange," JCAP **0903**, 018 (2009).

[11] K. Hinterbichler, L. Hui, and J. Khoury, "Conformal Symmetries of Adiabatic Modes in Cosmology," arXiv:1203.6351 [hep-th].

[12] P. Creminelli, J. Noreña, and M. Simonović, "Conformal Consistency Relations for Single-Field Inflation," arXiv:1203.4595 [hep-th].

[13] K. Hinterbichler, L. Hui, and J. Khoury, *to appear*.

[14] W. Goldberger, L. Hui, and A. Nicolis, *to appear*.

[15] T. Suyama and M. Yamaguchi, "Non-Gaussianity in the Modulated Reheating Scenario," Phys. Rev. D **77**, 023505 (2008).

[16] N. Sugiyama, E. Komatsu, and T. Futamase, "Non-Gaussianity Consistency Relation for Multi-Field Inflation," Phys. Rev. Lett. **106**, 251301 (2011).

[17] K. Smith, M. LoVerde, and M. Zaldarriaga, "A Universal Bound on N-Point Correlations from Inflation," Phys. Rev. Lett. **107**, 191301 (2011).

[18] T. Suyama, T. Takahashi, M. Yamaguchi, and S. Yokoyama, "On Classification of Models of Large Local-Type Non-Gaussianity," JCAP **1012**, 030 (2010).

[19] C. Byrnes, S. Nurmi, G. Tasinato, and D. Wands, "Inhomogeneous non-Gaussianity," JCAP **1203** 012 (2012).

[20] N. Sugiyama, "Consistency Relation for Multi-Field Inflation Scenario with All Loop

Contributions," arXiv:1201.4048 [gr-qc].

[21] C. Byrnes, M. Sasaki, and D. Wands, "The Primordial Trispectrum from Inflation," Phys. Rev. D **74**, 123519 (2006).

[22] D. Baumann, S. Ferrarro, D. Green, and K. Smith, *to appear.*

[23] D. Jeong and M. Kamionkowski, "Clustering Fossils from the Early Universe," arXiv:1203.0302 [astro-ph.CO].

[24] X. Chen and Y. Wang, "Quasi-Single-Field Inflation and Non-Gaussianities," JCAP **1004**, 027 (2010).

[25] D. Baumann and D. Green, "Signatures of Supersymmetry from the Early Universe," arXiv:1109.0292 [hep-th].

[26] S. Adler, "Consistency Conditions on the Strong Interactions Implied by a Partially Conserved Axial Vector Current," Phys. Rev. **137**, B1022 (1965).

[27] S. Weinberg, "Pion Scattering Lengths," Phys. Rev. Lett. **17**, 616 (1966).

[28] S. Weinberg, "Precise Relations between the Spectra of Vector and Axial Vector Mesons," Phys. Rev. Lett. **18**, 507 (1967).

[29] S. Weinberg, "The Quantum Theory of Fields. Vol. 2: Modern Applications," Cambridge, UK: Univ. Pr. (1996) 489 p.

[30] S. Coleman, "Aspects of Symmetry", Cambridge, UK: Univ. Pr. (1985).

[31] J. Donoghue, E. Golowich, and B. Holstein, "Dynamics of the Standard Model," Camb. Monogr. Part. Phys. Nucl. Phys. Cosmol. **2**, 1 (1992).

[32] E. Komatsu and D. Spergel, "Acoustic Signatures in the Primary Microwave Background Bispectrum," Phys. Rev. D **63**, 063002 (2001).

[33] C. Cheung, P. Creminelli, A. Fitzpatrick, J. Kaplan, and L. Senatore, "The Effective Field Theory of Inflation," JHEP **0803**, 014 (2008).

[34] L. Senatore and M. Zaldarriaga, "On Loops in Inflation," JHEP **1012**, 008 (2010);
L. Senatore and M. Zaldarriaga, "On Loops in Inflation II: IR Effects in Single Clock Inflation," arXiv:1203.6354 [hep-th];
G. Pimentel, L. Senatore, and M. Zaldarriaga, "On Loops in Inflation III: Time Independence of Zeta in Single Clock Inflation," arXiv:1203.6651 [hep-th].

[35] X. Chen, M. Huang, S. Kachru, and G. Shiu, "Observational Signatures and Non-Gaussianities of General Single-Field Inflation," JCAP **0701**, 002 (2007).

[36] R. Holman and A. Tolley, "Enhanced Non-Gaussianity from Excited Initial States," JCAP **0805**, 001 (2008).

[37] I. Agullo and L. Parker, "Non-Gaussianities and the Stimulated Creation of Quanta in the Inflationary Universe," Phys. Rev. D **83**, 063526 (2011).

[38] J. Ganc, "Calculating the Local-Type $f_{\rm NL}$ for Slow-Roll Inflation with a Non-Vacuum Initial State," Phys. Rev. D **84**, 063514 (2011).

[39] I. Agullo, J. Navarro-Salas, and L. Parker, "Enhanced Local-Type Inflationary Trispectrum from a Non-Vacuum Initial State," arXiv:1112.1581 [astro-ph.CO].

[40] S. Kundu, "Inflation with General Initial Conditions for Scalar Perturbations," JCAP **1202**, 005 (2012).

[41] V. Assassi, D. Baumann, and D. Green, *work in progress.*

[42] J. Maldacena and G. Pimentel, "On Graviton Non-Gaussianities during Inflation," JHEP **1109**, 045 (2011).

[43] P. Creminelli, "Conformal Invariance of Scalar Perturbations in Inflation," Phys. Rev. D **85**, 041302 (2012).

[44] A. Kehagias and A. Riotto, "Operator Product Expansion of Inflationary Correlators and Conformal Symmetry of de Sitter," arXiv:1205.1523 [hep-th].

[45] S. Weinberg, "Adiabatic Modes in Cosmology," Phys. Rev. D **67**, 123504 (2003).

[46] I. Low and A. Manohar, "Spontaneously Broken Spacetime Symmetries and Goldstone's Theorem," Phys. Rev. Lett. **88**, 101602 (2002).

[47] J. Bardeen, P. Steinhardt, and M. Turner, "Spontaneous Creation of Almost Scale-Free Density Perturbations in an Inflationary Universe," Phys. Rev. D **28**, 679 (1983).

[48] D. Salopek and J. Bond, "Non-Linear Evolution of Long-Wavelength Metric Fluctuations in Inflationary Models," Phys. Rev. D **42**, 3936 (1990).

[49] S. Weinberg, "Cosmology," Oxford, UK: Oxford Univ. Pr. (2008) 593 p.

[50] S. Weinberg, "The Quantum Theory of Fields. Vol. 1: Foundations," Cambridge, UK: Univ. Pr. (1995) 609 p.

[51] L. Senatore and M. Zaldarriaga, "A Note on the Consistency Condition of Primordial Fluctuations," arXiv:1203.6884 [astro-ph.CO].

[52] A. Nicolis and F. Piazza, "Spontaneous Symmetry Probing," arXiv:1112.5174 [hep-th].

[53] P. Creminelli, A. Nicolis, and E. Trincherini, "Galilean Genesis: An Alternative to Inflation," JCAP **1011**, 021 (2010).

[54] R. Flauger and E. Pajer, "Resonant Non-Gaussianity," JCAP **1101**, 017 (2011).

[55] X. Chen, "Primordial Non-Gaussianities from Inflation Models," Adv. Astron. **2010**, 638979 (2010).

[56] S. Weinberg, "Quantum Contributions to Cosmological Correlations," Phys. Rev. D **72**, 043514 (2005).

[57] D. Baumann and D. Green, "Equilateral Non-Gaussianity and New Physics on the Horizon," JCAP **1109**, 014 (2011).

[58] J. Ganc and E. Komatsu, "A New Method for Calculating the Primordial Bispectrum in the Squeezed Limit," arXiv:1006.5457, [astro-ph.CO].

[59] J. Collins, D. Soper, and G. Sterman, "Factorization of Hard Processes in QCD," Adv. Ser. Direct. High Energy Phys. **5**, 1 (1988).

[60] PLANCK, `http://www.rssd.esa.int/index.php?project=Planck`

[61] SDSS, `http://www.sdss3.org/`

[62] F. Schmidt and M. Kamionkowski, "Halo Clustering with Non-Local Non-Gaussianity," Phys. Rev. D **82**, 103002 (2010).

[63] EUCLID, `http://www.euclid-ec.org/`

[64] LSST, `http://www.lsst.org/`

[65] PIXIE, A. Kogut *et al.*, "The Primordial Inflation Explorer (PIXIE): A Nulling Polarimeter for Cosmic Microwave Background Observations," JCAP **1107**, 025 (2011).

[66] CMBPol, D. Baumann *et al.* [CMBPol Study Team Collaboration], "CMBPol Mission Concept Study: Probing Inflation with CMB Polarization," AIP Conf. Proc. **1141**, 10 (2009).

[67] P. Creminelli, S. Dubovsky, A. Nicolis, L. Senatore, and M. Zaldarriaga, "The Phase Transition to Slow-Roll Eternal Inflation," JHEP **0809**, 036 (2008).